

Carbon-rich RR Lyr type stars.

George Wallerstein ¹

Department of Astronomy, University of Washington, Seattle, WA 98195,
wall@astro.washington.edu

V. V. Kovtyukh² and S. M. Andrievsky²

Department of Astronomy and Astronomical Observatory, Odessa National University,
T.G. Shevchenko Park, 65014, Odessa, Ukraine

val@deneb1.odessa.ua, scan@deneb1.odessa.ua

Received _____; accepted _____

ABSTRACT

We have derived CNO abundances in 12 RR Lyrae stars. Four stars show $[C/Fe]$ near 0.0 and two stars show $[C/Fe] = 0.52$ and 0.65 . Red giant branch stars, which are known to be the predecessors of RR Lyrae stars, generally show a deficiency of carbon due to proton captures during their evolution from the main sequence up the giant branch. We suggest that the enhancement of carbon is due to production during the helium flash combined with mixing to the surface by vigorous convection induced by the flash itself.

Subject headings: stars: RR Lyr type, individual: RS Boo, SW And, AN Ser, RR Lyr, UY Boo, SU Dra, VY Ser, OX Her, V486 Her, XZ Cet, KP Cyg, UY CrB

1. Introduction

It is generally agreed that the RR Lyrae stars have evolved from the red giant branch (RGB) to their present position on the horizontal branch (HB) of globular clusters when the helium flash initiates the triple-alpha reaction in the degenerate core. Because of the degeneracy, combined with the great temperature dependence of the triple-alpha reaction, the core of the red giant heats extremely rapidly leading to very vigorous convection that must be treated by hydrodynamic methods rather than the conventional theory of steady convection. Recent calculations have been described by Dearborn, Lattanzio & Eggleton (2006) as well as by Mocak et al. (2008). There remain many uncertainties in the calculations but it is generally agreed that the star is not disrupted. This is demonstrated by the very existence of HB stars. At the same time the degree of mixing remains uncertain with the distinct possibility that freshly minted ^{12}C may reach the surface. In addition the ^{12}C must pass through a region rich in hydrogen so it may be reprocessed to ^{13}C and ^{14}N .

As part of a small survey of the chemical composition of specially selected RR Lyrae stars and comparison stars of known metallicity we have found at least two carbon-rich objects.

2. Observations

The spectra of our program stars were obtained with the echelle spectrograph of the Apache Point Observatory (APO). By using a prism as cross-disperser the APO echelle covers all wavelengths from 3500 Å to 10400 Å. However the red-sensitive 2048×2048 chip has decreasing sensitivity below 4000 Å and beyond 9000 Å. The resolving power is about 35000. Exposure times were usually about 20–30 minutes. We estimated the S/N ratio at the continuum level depending upon the wavelength interval to be about 70–150 per pixel

after combining multiple exposures that had been taken sequentially. Table 1 contains the dates, JD, phases, and derived values of T_{eff} and $\log g$ for each phase.

3. Data Reduction and Analysis

3.1. Data Reduction

The program spectra were extracted from the raw frames using standard IRAF procedures. The continuum level placement, wavelength calibration and equivalent width measurements were performed with DECH20 code (Galazutdinov 1992).

3.2. Atmospheric Parameters and Abundance Analysis

The elemental abundances were derived using the Kurucz’s WIDTH9 code with atmosphere models interpolated from the ATLAS9 model grid. We used $\log gf$ values derived from an inverted solar analysis (Kovtyukh & Andrievsky 1999).

Atmosphere parameters (T_{eff} , $\log g(\text{sp})$, V_t) were derived by enforcing traditional spectroscopic criteria. Lines of Fe I were forced to yield zero slope in the relations between total iron abundance and excitation potential. The total abundances of iron as predicted from Fe I and Fe II lines were equalized by adjusting the model gravity (i.e. to yield a ”spectroscopic gravity”).

4. CNO abundances

To derive the carbon abundance the following set of the lines was used: 6001.12 Å, 6010.68 Å, 6014.83 Å, 6587.61 Å, 6655.51 Å, 7085.47 Å, 7087.83 Å, 7111.48 Å, 7113.18 Å,

Table 1: Observations of stars and their fundamental characteristics

Star	Period (days)	Date	JD 2450000+	ϕ	T _{eff} K	log g	V _t km s ⁻¹	Remarks
RS Boo	0.377	2002-05-23	2417.628	.440(b)	6500	2.9	3.3	
		2006-03-21	3816.745	.311(b)	6700	3.5	3.5	
SW And	0.442	2005-01-15	3386.542	.797(c)	6200	2.5	3.8	
AN Ser	0.522	2002-05-23	2417.664	.886(f)	7500	2.8	4.8	
		2002-06-22	2447.889	.780(f)	6150	2.4	3.4	
		2002-06-23	2448.638	.214(f)	6300	2.6	3.7	
RR Lyr	0.567	2003-06-14	2805.790	.268(e)	6200	2.6	3.8	
		2005-07-05	3556.950	.448(e)	6100	2.8	2.3	
UY Boo	0.651	2002-05-23	2417.733		6200	2.0	2.8	Na D1,D2 double
SU Dra	0.660	2006-03-21	3816.731	.676(c)	6100	2.5	3.5	H α emission or double
VY Ser	0.714	2002-03-02	2336.876	.000(e)	6800	2.0	3.0	H α emission or double
		2002-05-22	2417.688	.168(e)	6350	2.0	3.0	
		2002-06-21	2447.637	.107(e)	6450	2.2	3.0	
		2002-06-22	2448.689	.581(e)	6000	2.3	3.7	
OX Her	0.757	2005-06-18	3539.757	.756(i)	6000	2.7	3.3	
		2005-06-20	3541.702	.324(i)	6800	3.0	3.8	
V486 Her	0.806	2004-06-05	3162.756	.260(d)	6700	3.4	3.8	H α emission or double
		2005-03-26	3456.880	.210(d)	6700	3.3	3.5	H α emission or double
		2005-06-17	3539.699	.972(d)	7000	3.8	4.5	H α blue-shifted emission
		2005-06-19	3541.762	.532(d)	6200	3.2	4.0	
XZ Cet	0.823(a)	2003-01-24	2664.596	.167(h)	6400	2.3	3.0	
		2003-02-07	2678.562	.134(h)	6450	2.2	2.8	
KP Cyg	0.856	2005-11-13	3687.606	.135(d)	7050	2.6	2.5	
		2006-11-05	4044.672	.299(d)	6600	3.0	3.3	
		2007-05-01	4221.963	.430(d)	6450	3.4	3.5	H α emission
		2005-09-24	3637.607	.720(d)	6300	2.4	3.2	H α emission
		2006-10-03	4011.720	.800(d)	6650	3.0	4.2	H α emission
		2005-09-13	3626.711	.991(d)	7400	3.0	2.5	
UY CrB	0.929	2004-06-05	3162.681	.491(g)	6150	2.3	3.8	
		2005-03-26	3456.989	.245(g)	6700	2.4	3.4	
		2005-09-13	3626.644	.839(g)	6300	2.9	3.0	

Remark: Phases are from:

a - XZ Cet was classified as an anomalous cepheid according to Teays & Simon (1985), but that classification is uncertain. b - Jones R.V. et al., 1987, ApJ 314, 605 c - Liu T.& Janes K.A., 1990, ApJ 354, 273 d - Loomis Ch., Schmidt E.G., Simon N.R., 1988, MNRAS 235, 1059 e - Takeda Y., Honda S., Aoki W., Takada-Hidai M., Zhao G., Chen Yu-Qin, Shi Jian-Rong, 2006, PASJ 58, 389 f - Lub J., 1977, A&AS 29, 345 g - Schmidt E.G., 2002, AJ 123, 965 h - Schmidt E.G., 2002, AJ 123, 965 i - GCVS, 1984, 1985

7115.19 Å, 7116.99 Å, 8335.15 Å. The Nitrogen abundance was found from the lines 7442.29 Å, 7468.30 Å, 8216.34 Å, 8242.39 Å, 8629.16 Å, 8683.39 Å, 8703.25 Å, 8711.67 Å, 8718.76 Å. These lines are seen only in the stars with a quite high metallicity. For the metal-poor ones only some of them were present. The abundance of oxygen in the program stars was derived from the following lines: 6156.77 Å, 6158.18 Å, and 6300.30 Å. Equivalent widths will be presented in a full paper on abundances of all available elements in the stars in Table 1.

As an example, in Fig. 1 we show spectra in the region of the C I lines near 7110 Å. Note that all stars showing these lines were observed near the same phase, 0.5, and hence have very nearly the same temperatures.

The results for the CNO abundances in our program stars are given in Table 2 (we have shown the derived abundances on the usual scale letting $\log N(H)=12.0$).

Only a few comparisons may be made with other analyses of RR Lyr star abundances. For SW And Clementini et al. (1995) found $[C/Fe] = -0.10$ while we found $+0.09$. The difference is almost entirely due to their solar abundance of C of 8.56 (Anders and Grevesse 1990) and our more recent value of 8.39 (Asplund et al. 2005). For $[O/Fe]$ they found -0.03 and we found $+0.23$ which corresponds closely to the revision of the solar oxygen abundance from 8.93 to 8.66.

It is also worth noting the carbon and s-process enhancements found by Preston et al. (2006) in TY Gru, due probably to mass transfer from a companion. We have analysed the heavy element content of the two C-rich stars, KP Cyg and UY CrB (to be reported on later) and found no significant enhancements.




Fig. 1.— C I lines in the program stars. Note that all stars were observed at nearly the same phase so their effective temperatures are similar.

5. Discussion

It is well known that in evolving metal-poor stars the first dredge-up depletes C in the stellar atmosphere, and that the additional mixing that occurs at the red giant clump further depletes it. Normal or excess carbon indicates that additional carbon has been injected into the atmosphere from deeper layers, presumably from the helium flash without being reprocessed to N by proton capture on the way up. Two stars, KP Cyg and UY CrB, where the carbon excess in units of $[C/Fe]$ was found to be 0.52 and 0.65, have rather long periods and $[Fe/H]$ values that are relatively high.

Almost all stars in which it was possible to determine the N abundance show an excess of that element. Since there is no significant carbon deficiency, the excess of N must be left over from the first and red giant clump mixing events. Three stars with $[Fe/H]$ near zero show no N excess.

Only for six stars were we able to measure the oxygen abundance. Except in UY CrB, there is no indication of an oxygen excess having been generated during the helium flash by the $^{12}C (\alpha, \gamma)^{16}O$.

Some comments on individual stars are worthwhile. KP Cyg appears to be one of the most metal-rich RR Lyrae stars known. Its carbon abundance is 0.7 dex greater than solar but its oxygen abundance is not enhanced. Since its $[Fe/H]$ value is -0.4 UY CrB probably started with $\log N(C) = 8.0$. It now shows $\log N(C)=8.6$. Its $\log N(O)$ must have started near 8.26 and now has reached 8.85 indicating additional processing of ^{12}C to ^{16}O by alpha-capture. For both stars nitrogen is substantially enhanced as might be expected as carbon-rich material is subject to proton capture on its way to the surface. For the metal-poor star, RR Lyr, with $[Fe/H] = -1.2$ the original carbon abundance was probably about $\log N(C) = 7.2$ which is likely to have been depleted to 6.7-7.0 at the RG tip and is 7.4 at present. Its present $\log N(N)=7.4$ which must also be an enhancement from a

presumed initial $\log N(N) = 6.6$.

To compare the RR Lyrae stars with less evolved objects whose origin is likely to be similar we show abundances in various RGB stars in Table 3. For this we have used data for stars in M71, M4 and 47 Tuc all of which have metallicities similar to those in the C-rich RR Lyrae stars.

It appears that some RR Lyrae stars show a significant enhancement in carbon as compared to red giants of comparable metallicity. Apparently, an increased carbon abundance at the evolutionary stage after the red giant phase is caused by the deep convective mixing. Calculations by Fujimoto et al.(2000) and by Schlattl et al.(2002) for the He-flash in extremely metal-poor stars show that thermally driven convection in the core can reach the H-burning shell. When H mixes with the freshly synthesized carbon rapid proton capture can release further energy to drive the convection to the outer layers of the star.

We are currently expanding our high resolution survey of RR Lyrae stars to establish any correlation of carbon excesses with period, metallicity or other parameters.

SMA and VVK would like to express their gratitude for the Kenilworth Fund of the New York Community Trust for the financial support of this study. The individual financial support from Kenilworth Fund was made possible through CRDF.

We thank Marta Mottini and Wenjin Huang for reading the manuscript and making some good suggestions.

The search of the information about the program stars was made using SIMBAD.

Table 2: CNO abundances

Star	logN(C)	[C/Fe]	logN(N)	[N/Fe]	logN(O)	[O/Fe]	[Fe/H]
RS Boo	8.34	−0.06	8.00	+0.21	8.44	−0.21	+0.01
SW And	8.19	+0.09	7.69	+0.20	8.60	+0.23	−0.29
AN Ser	8.37	+0.00	7.92	+0.12	8.71	+0.03	+0.02
RR Lyr	6.66	−0.56	<7.50	<+0.89	8.30	−0.19	−1.17
UY Boo	<6.10	<+0.44	<7.60	<+2.55	<8.30	<+2.37	−2.73
SU Dra	<6.20	<−0.25	<7.20	<+0.36	<8.50	<+1.78	−1.94
VY Ser	<6.20	<−0.53	<7.20	<+1.08	<8.40	<+1.40	−1.66
OX Her	<6.50	<−0.39	<7.20	<+0.92	<8.50	<+1.34	−1.50
V486 Her	<6.90	<−0.60	<8.00	<+1.11	<8.30	<+0.56	−0.89
XZ Cet	<6.10	<−0.51	<7.10	<+1.10	<8.50	<+1.62	−1.78
KP Cyg	9.09	+0.52	8.86	+0.90	8.77	−0.07	+0.18
UY CrB	8.64	+0.65	8.64	+1.26	8.85	+0.59	−0.40
Sun	8.39		7.78		8.66		0.00

For the Sun we adopted the CNO abundances as recommended by Asplund et al. (2004), Asplund et al. (2005ab).

For UY Boo, SU Dra, VY Ser, OX Her, V486 Her and XZ Cet only upper limit of CNO abundances is given. For the nitrogen abundance in RR Lyr also the upper limit is indicated.

Table 3: CNO abundances in globular clusters

Cluster	Type of star	[Fe/H]	logN(C)	logN(N)	logN(O)	$^{12}\text{C}/^{13}\text{C}$	references
M 71	CNweak	−0.79	7.07	7.71	8.24	8.3	a
M 71	CNstrong	−0.78	7.11	8.07	8.08	6.0	a
M 71	AGB	−0.89	6.76	8.04	8.14	5	a
47 Tuc	GB	−0.77	7.48:	8.29:	8.35	4–12	b,d
M 4	GB	−1.34	6.97:	7.83:	7.87:	4–8	b
M 4	AGB	−1.18	6.64	7.35	7.73	4.5	c

References: a - Briley et al. (1997), b - Brown & Wallerstein (1990), c - Ivans, I.I., et al. (1999), d- Alves-Brito, et al. (2005).

REFERENCES

- Alves-Brito, A. et al. 2005 A&A 435, 657
- Anders E., Grevesse N., 1990 Geochim. Cosmochim. Acta 53,197
- Asplund M., Grevesse N., Sauval A.J., Allende Prieto C., Kiselman D., 2004, A&A 417, 751
- Asplund M., Grevesse N., Sauval A.J., Allende Prieto C., Blomme R., 2005, A&A 431, 693
- Asplund M., Grevesse N., Sauval A.J., 2005, ASP Conference Series 336, 25
- Briley M.M., Smith V.V., Kink J., Lambert D.L., 1997, AJ 113, 306
- Brown J.A., Wallerstein G. 1992, AJ 100,1561
- Clementini G., Carretta E., Gratton R., Merighi R., Mould J.R., McCarthy J.K., 1995, AJ 110, 2319
- Fujimoto M.Y., Ikeda Y., Iben, I.Jr., 2000, ApJ 529, 25
- Galazutdinov G.A., 1992, Preprint SAO RAS, 92
- Ivans, I.I. et al. 1999 AJ 118, 1273
- Dearborn D.S.P., Lattanzio J.C., Eggleton P.P., 2006 ApJ 639, 4
- Kovtyukh V.V., Andrievsky S.M., 1999, A&A 351, 597
- Kurucz R. 1996, In Model Atmospheres and Spectrum Synthesis, ASPC 108, 270
- Mocak M., Mueller E., Weiss A., Kifonides K., 2008, A&A, in press
- Schlattl H., Salaris M., Cassisi S., Weiss A., 2002 A&A, 395, 77
- Teays T.J., Simon N.R., 1985, ApJ 290, 683

